



Hai Nguyen<sup>1</sup> Matthias Katzfuss<sup>2</sup> Noel Cressie<sup>3</sup>  
Amy Braverman<sup>1</sup>

<sup>3</sup>Department of Statistics,  
The Ohio State University



National Aeronautics and  
Space Administration

**Jet Propulsion Laboratory**  
California Institute of Technology  
Pasadena, California

## Outline

### Introduction

Prelude: spatial (non-temporal) interpolation for massive datasets

Spatio-Temporal Data Fusion

Application to Lower Atmosphere CO<sub>2</sub>

Conclusion



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Definition

*Data fusion is the process of combining information from heterogeneous sources into a single composite picture of the relevant process, such that the composite picture is generally more accurate and complete than that derived from any single source alone (Hall, 2004).*



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Motivation: remote sensing data

### What is the benefit of data fusion?

- ▶ Remote sensing data are often incomplete, sparse, and spatially and temporally heterogeneous. Our goal is to infer the true physical process from all available data sources.
- ▶ Data fusion capitalizes on complementary strengths of the individual datasets to minimize prediction errors.
- ▶ Correlation in *space* and *time* can be exploited for improved accuracy.





National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Motivating example

### Estimating lower atmosphere CO<sub>2</sub>

- ▶ The lower atmosphere (below 500 hPa )is where CO<sub>2</sub> enters and exits the atmosphere. This may be a proxy for 'sources' and 'sinks'.
- ▶ No satellite instrument currently provides measurements of CO<sub>2</sub> globally near the Earth's surface.
- ▶ The Greenhouse gases Observing SATellite (GOSAT) provides total-column CO<sub>2</sub>, while the Atmospheric InfraRed Sounder (AIRS) provides mid-tropospheric CO<sub>2</sub>.
- ▶ Approximations to lower-atmospheric CO<sub>2</sub> may be made by deriving joint predictions of total-column CO<sub>2</sub> and mid-tropospheric CO<sub>2</sub> and taking a (weighted) difference.



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Example illustration

### Example of satellite orbits and footprints




Figure: Example of different footprints ( Source: Amy Braverman)



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Difficulties of data fusion

Difficulties encountered when fusing remote sensing datasets:

- ▶ Massive size,
- ▶ Change of support,
- ▶ Isotropy and stationarity,
- ▶ Accounting for instruments' biases.



National Aeronautics and  
Space Administration

**Jet Propulsion Laboratory**  
California Institute of Technology  
Pasadena, California

## Outline

Introduction

Prelude: spatial (non-temporal) interpolation for massive datasets

Spatio-Temporal Data Fusion

Application to Lower Atmosphere CO<sub>2</sub>

Conclusion



National Aeronautics and  
Space Administration

**Jet Propulsion Laboratory**  
California Institute of Technology  
Pasadena, California

## Motivation

A special case of Spatio-Temporal Data Fusion is non-temporal spatial interpolation. It is convenient to introduce the ideas of STDF by talking about the spatial-only case first.



We assume the data from an instrument is generated according to the following model:

$$\mathbf{Z} = (Z(B_1), Z(B_2), \dots, Z(B_N))',$$
$$Z(B_j) = \frac{1}{|D \cap B_j|} \left\{ \sum_{\mathbf{u} \in D \cap B_j} Y(\mathbf{u}) \right\} + \epsilon(B_j); \quad B_j \subset \mathbb{R}^d,$$

where

- ▶  $D$  is a discretized domain made up of Basic Areal Units (BAU),
- ▶  $B_{ij}$  is the  $j$ th footprint from dataset  $i$  ( $i = 1, 2$ ),
- ▶  $\mathbf{Z}_i$  is the vector of response variable from dataset  $i$ ,
- ▶  $Y(\cdot)$  is the true process,
- ▶  $\epsilon_i(B_{ij})$  is the error process.



We assume that the spatial process has the following linear mixed model (Cressie and Johannesson, 2008),

$$Y(\mathbf{s}) = \mathbf{t}(\mathbf{s})'\boldsymbol{\alpha} + \mathbf{S}(\mathbf{s})'\boldsymbol{\eta} + \xi(\mathbf{s}),$$

where

- ▶  $\xi(\cdot)$  is a fine-scale variation process (white noise) w/ variance  $\sigma_\xi^2$ ,
- ▶  $\mathbf{t}(\mathbf{s})'\boldsymbol{\alpha}$  accounts for a linear trend,
- ▶  $\boldsymbol{\eta}$  is an  $r$ -dimensional Gaussian random vector  $\text{var}(\boldsymbol{\eta})$ ,
- ▶  $\mathbf{S}(\mathbf{s})$  is an  $r$ -dimensional spatial basis expansion of  $\mathbf{s}$ .



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Covariance model

Given the linear mixed model, the covariance model is,

$$\mathbf{\Sigma} \equiv \text{var}(\mathbf{Z}) = \mathbf{S}'\mathbf{K}\mathbf{S} + \sigma_{\xi}^2\mathbf{E} + \sigma_{\epsilon}^2\mathbf{V}.$$

where

- ▶  $\mathbf{K} = \text{var}(\boldsymbol{\eta})$ : fixed dimension  $r \times r$ ,
- ▶  $\mathbf{S} \equiv (\mathbf{S}(B_1), \dots, \mathbf{S}(B_N))'$ ,
- ▶  $\mathbf{E}$  and  $\mathbf{V}$  are known matrices.





National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Prediction form

The optimal (linear unbiased) predictor of  $Y(\mathbf{s})$  can be written as

$$\hat{Y}(\mathbf{s}) = \mathbf{a}'\mathbf{Z},$$

where  $\mathbf{a}$  is an  $N$ -dimensional vector of kriging coefficients.



We wish to minimize,

$$\begin{aligned} E(Y(\mathbf{s}) - \hat{Y}(\mathbf{s}))^2 &= \text{var}(Y(\mathbf{s}) - \mathbf{a}'\mathbf{Z}), \\ &= \text{var}(Y(\mathbf{s})) - 2\mathbf{a}'\text{cov}(\mathbf{Z}, Y(\mathbf{s})) + \mathbf{a}'\text{var}(\mathbf{Z})\mathbf{a}, \end{aligned}$$

with respect to  $\mathbf{a}$ , subject to the unbiasedness constraint,

$$\mathbf{0} = \mathbf{a}'\mathbf{T} - \mathbf{t}(\mathbf{s})'.$$

Solving using the method of Lagrange multipliers, the optimal kriging coefficients  $\mathbf{a}$  is,

$$\mathbf{a}' = (\mathbf{c}' + (\mathbf{t}(\mathbf{s})' - \mathbf{c}'\mathbf{\Sigma}^{-1}\mathbf{T})(\mathbf{T}'\mathbf{\Sigma}^{-1}\mathbf{T})^{-1}\mathbf{T}')\mathbf{\Sigma}^{-1}.$$



## Prediction and standard error equations

We can interpolate at a new location with the following,

$$\begin{aligned}Y(\mathbf{s})^{FRK} &= \mathbf{a}' \mathbf{Z}, \\ \sigma(\mathbf{s})^{SSDF} &= \left( E(Y(\mathbf{s})^{SSDF} - Y(\mathbf{s}))^2 \right)^{\frac{1}{2}} \\ &= \left( \mathbf{S}(\mathbf{s})' \mathbf{K} \mathbf{S}(\mathbf{s}) + \sigma_{\xi}^2 - 2\mathbf{a}' (\mathbf{S}' \mathbf{K} \mathbf{S}(\mathbf{s}) + \mathbf{b}(\mathbf{s})) \right. \\ &\quad \left. + \mathbf{a}' (\mathbf{S}' \mathbf{K} \mathbf{S} + \sigma_{\xi}^2 \mathbf{E} + \mathbf{V}) \mathbf{a} \right)^{\frac{1}{2}},\end{aligned}$$

where

$$\mathbf{b}(\mathbf{s}) = \text{cov}(\xi, \xi(\mathbf{s})).$$

This is called **Fixed Rank Kriging**.



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Advantages of FRK

- ▶ Inversion of  $\Sigma$  is computationally scalable using the Sherman-Morrison-Woodbury formula (Henderson and Searle, 1981),

$$\Sigma^{-1} = \mathbf{U}^{-1} - \mathbf{U}^{-1}\mathbf{S}'(\mathbf{K}^{-1} + \mathbf{S}\mathbf{U}^{-1}\mathbf{S}')^{-1}\mathbf{S}\mathbf{U}^{-1},$$

where  $\mathbf{U} = \sigma_{\xi}^2 \mathbf{E} + \mathbf{V}$ .

- ▶ No assumption of isotropy or stationarity.
- ▶ Handles change of support.
- ▶ Able to handle known systematic instrument biases.



National Aeronautics and  
Space Administration

**Jet Propulsion Laboratory**  
California Institute of Technology  
Pasadena, California

## Outline

### Introduction

Prelude: spatial (non-temporal) interpolation for massive datasets

### Spatio-Temporal Data Fusion

Application to Lower Atmosphere CO<sub>2</sub>

### Conclusion



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Extending FRK to spatio-temporal data fusion

We develop an extension of FRK called Spatio-Temporal Data Fusion, which has the following properties,

- ▶ Ability to derive joint-estimates of two or more processes,
- ▶ Ability to exploit both spatial and temporal dependence in the data.

Main ideas behind STDF

- ▶ We account for temporal dependence using a first-order auto-regressive model for  $\eta$ .
- ▶ We do optimal predictions using a variant of the Kalman smoother.



We assume the data from an instrument is generated according to the following model:

$$Z_t^{(k)}(A) = \frac{1}{|D \cap A|} \left\{ \sum_{\mathbf{s} \in D \cap A} Y_t^{(k)}(\mathbf{s}) \right\} + \epsilon_t^{(k)}(A),$$

where

- ▶  $A \subset R^d$ ,  $k = 1, 2$ ,  $t = 1, 2, \dots, T$ ,
- ▶  $Y_t^{(k)}(\cdot)$  is the true process,
- ▶  $\epsilon_t^{(k)}(A)$  is the error process.



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Process model

We assume that the  $k$ -th process has the following form,

$$Y_t^{(k)}(\mathbf{s}) = \mathbf{x}_t^{(k)}(\mathbf{s})' \boldsymbol{\alpha}_t^{(k)} + \mathbf{S}_t^{(k)}(\mathbf{s})' \boldsymbol{\eta}_t^{(k)} + \xi_t^{(k)}(\mathbf{s}); \quad \mathbf{s} \in D.$$

where  $\mathbf{x}_t^{(k)}(\cdot)$ ,  $\boldsymbol{\alpha}_t^{(k)}$ ,  $\mathbf{S}_t^{(k)}(\cdot)$ ,  $\boldsymbol{\eta}_t^{(k)}$ , and  $\xi_t^{(k)}(\cdot)$  are defined in an analogous fashion to the corresponding spatial-only terms.





National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Instrument bias

To allow for bias, we assume that the measurement-error process may have non-zero mean,

$$\begin{aligned} E(\epsilon_t^{(k)}(A)) &= c^{(k)} E(Y^{(k)}(A)) \\ &= c^{(k)} \mathbf{x}(A)' \boldsymbol{\alpha}. \end{aligned}$$

- The multiplicative bias coefficients  $\{c^{(k)} : k = 1, 2\}$  are assumed known.



At time  $t$ , we can stack datasets  $\mathbf{Z}_t^{(1)}$  and  $\mathbf{Z}_t^{(2)}$  to form a joint vector,

$$\begin{pmatrix} \mathbf{Z}_t^{(1)} \\ \mathbf{Z}_t^{(2)} \end{pmatrix} = \begin{pmatrix} \mathbf{X}_t^{(1)} & \mathbf{0} \\ \mathbf{0} & \mathbf{X}_t^{(2)} \end{pmatrix} \begin{pmatrix} \boldsymbol{\alpha}_t^{(1)} \\ \boldsymbol{\alpha}_t^{(2)} \end{pmatrix} + \begin{pmatrix} \mathbf{S}_t^{(1)} & \mathbf{0} \\ \mathbf{0} & \mathbf{S}_t^{(2)} \end{pmatrix} \begin{pmatrix} \boldsymbol{\eta}_t^{(1)} \\ \boldsymbol{\eta}_t^{(2)} \end{pmatrix} \\ + \begin{pmatrix} \boldsymbol{\xi}_t^{(1)} \\ \boldsymbol{\xi}_t^{(2)} \end{pmatrix} + \begin{pmatrix} \boldsymbol{\epsilon}_t^{(1)} \\ \boldsymbol{\epsilon}_t^{(2)} \end{pmatrix},$$

or equivalently,

$$\mathbf{Z}_t = \mathbf{X}_t \boldsymbol{\alpha}_t + \mathbf{S}_t \boldsymbol{\eta}_t + \boldsymbol{\xi}_t + \boldsymbol{\epsilon}_t.$$



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## First-order auto-regressive temporal model

We assume that the covariance parameter  $\eta_t$  evolves according to a first-order auto regressive model,

$$\eta_t | \eta_{t-1}, \dots, \eta_0 \sim N_r(\mathbf{H}_t \eta_{t-1}, \mathbf{U}_t); \quad t = 1, 2, \dots,$$

where

- ▶ The initial state is  $\eta_0 \sim N_r(\mathbf{0}, \mathbf{K}_0)$ ,
- ▶ The matrices  $\mathbf{H}_t$  and  $\mathbf{U}_t$  are called the propagator matrix and the innovation matrix.



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Spatial-Temporal Data Fusion

Given data from two different instruments  $\mathbf{Z}_1, \dots, \mathbf{Z}_T$ , we wish to optimally estimate the true processes at a set of locations  $P \subset D$  at time  $t \in \{1, \dots, T\}$ .

Assuming that the parameters  $\alpha$ ,  $\mathbf{K}_0$ ,  $\{(\sigma_\epsilon^{(k)})^2\}$ ,  $\{(\sigma_\xi^{(k)})^2\}$ ,  $\mathbf{H}_t$ , and  $\mathbf{U}_t$  are known, we can optimally estimate the posterior expectations and covariances for  $\{\eta_t\}$  and  $\{\xi_t^P\}$  using a variant of Kalman smoothing.



## Kalman smoothing

Let  $\mathbf{Z}_{1:\tilde{t}} \equiv (\mathbf{Z}'_1, \dots, \mathbf{Z}'_{\tilde{t}})'$ , we define,

- ▶  $\boldsymbol{\eta}_{t|\tilde{t}} \equiv \mathbb{E}(\boldsymbol{\eta}_t | \mathbf{Z}_{1:\tilde{t}}),$
- ▶  $\boldsymbol{\xi}_{t|\tilde{t}}^P \equiv \mathbb{E}(\boldsymbol{\xi}_t^P | \mathbf{Z}_{1:\tilde{t}}),$
- ▶  $\mathbf{P}_{t|\tilde{t}} \equiv \text{var}(\boldsymbol{\eta}_t | \mathbf{Z}_{1:\tilde{t}}),$
- ▶  $\mathbf{R}_{t|\tilde{t}}^P \equiv \text{var}(\boldsymbol{\xi}_t^P | \mathbf{Z}_{1:\tilde{t}}),$
- ▶  $\mathbf{W}_{t:\tilde{t}}^P \equiv \text{cov}(\boldsymbol{\eta}_t, \boldsymbol{\xi}_t^P | \mathbf{Z}_{1:\tilde{t}}).$



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Kalman smoothing

We first initialize  $\boldsymbol{\eta}_{0|0} = \mathbf{0}$  and  $\mathbf{P}_{0|0} = \mathbf{K}_0$ . The one-step ahead forecasts are,

$$\begin{aligned}\boldsymbol{\eta}_{t|t-1} &= \mathbf{H}_t \boldsymbol{\eta}_{t-1|t-1} \\ \mathbf{P}_{t|t-1} &= \mathbf{H}_t \mathbf{P}_{t-1|t-1} \mathbf{H}_t' + \mathbf{Q}_t.\end{aligned}$$



The filtering quantities for  $t = 1, \dots, T$  are:

$$\eta_{t|t} = \eta_{t|t-1} + \mathbf{P}_{t|t-1} \mathbf{S}_t' [\mathbf{S}_t \mathbf{P}_{t|t-1} \mathbf{S}_t' + \mathbf{D}_t]^{-1} (\mathbf{Z}_t - \mathbf{Q} \mathbf{X}_t \alpha_t - \mathbf{S}_t \eta_{t|t-1})$$

$$\xi_{t|t}^P = \mathbf{C}_t^{PZ} \mathbf{E}_t^{PZ} [\mathbf{S}_t \mathbf{P}_{t|t-1} \mathbf{S}_t' + \mathbf{D}_t]^{-1} (\mathbf{Z}_t - \mathbf{Q} \mathbf{X}_t \alpha_t - \mathbf{S}_t \eta_{t|t-1})$$

$$\mathbf{P}_{t|t} = \mathbf{P}_{t|t-1} - \mathbf{P}_{t|t-1} \mathbf{S}_t' [\mathbf{S}_t \mathbf{P}_{t|t-1} \mathbf{S}_t' + \mathbf{D}_t]^{-1} \mathbf{S}_t \mathbf{P}_{t|t-1}$$

$$\mathbf{R}_{t|t}^P = \mathbf{C}_t^P \mathbf{E}_t^P - \mathbf{C}_t^{PZ} \mathbf{E}_t^{PZ} [\mathbf{S}_t \mathbf{P}_{t|t-1} \mathbf{S}_t' + \mathbf{D}_t]^{-1} (\mathbf{E}_t^{PZ})' (\mathbf{C}_t^{PZ})',$$

$$\mathbf{W}_{t|t}^P = -\mathbf{P}_{t|t-1} \mathbf{S}_t' [\mathbf{S}_t \mathbf{P}_{t|t-1} \mathbf{S}_t' + \mathbf{D}_t]^{-1} (\mathbf{E}_t^{PZ})' (\mathbf{C}_t^{PZ})',$$

where  $\text{var}(\xi_t^P) = \mathbf{C}_t^P \mathbf{E}_t^P$ ,  $\text{cov}(\xi_t^P, \xi_t) = \mathbf{C}_t^{PZ} \mathbf{E}_t^{PZ}$ , and  $\mathbf{Q}$  is a diagonal matrix with  $\{c^{(k)}\}$  along the diagonal.



We obtain the smoothing quantities by updating “backwards” in time (i.e., for  $t = T - 1, T - 2, \dots, 0$ ):

$$\begin{aligned}\eta_{t|T} &= \eta_{t|t} + \mathbf{J}_t(\eta_{t+1|T} - \eta_{t+1|t}) \\ \xi_{t|T}^P &= \xi_{t|t}^P + \mathbf{B}_t(\eta_{t+1|T} - \eta_{t+1|t}) \\ \mathbf{P}_{t|T} &= \mathbf{P}_{t|t} + \mathbf{J}_t(\mathbf{P}_{t+1|T} - \mathbf{P}_{t+1|t})\mathbf{J}_t' \\ \mathbf{R}_{t|T}^P &= \mathbf{R}_{t|t}^P + \mathbf{B}_t(\mathbf{P}_{t+1|T} - \mathbf{P}_{t+1|t})\mathbf{B}_t' \\ \mathbf{W}_{t|T}^P &= \mathbf{W}_{t|t}^P + \mathbf{J}_t(\mathbf{P}_{t+1|T} - \mathbf{P}_{t+1|t})\mathbf{B}_t'\end{aligned}$$

where

$$\begin{aligned}\mathbf{J}_t &\equiv \mathbf{P}_{t|t}\mathbf{H}_{t+1}'\mathbf{P}_{t+1|t}^{-1} \\ \mathbf{B}_t &\equiv -\mathbf{C}_t^{PZ}\mathbf{E}_t^{PZ} [\mathbf{S}_t\mathbf{P}_{t|t-1}\mathbf{S}_t' + \mathbf{D}_t]^{-1}\mathbf{S}_t\mathbf{P}_{t|t-1}\mathbf{H}_{t+1}'\mathbf{P}_{t+1|t}^{-1}.\end{aligned}$$





National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Prediction equation

Having obtained the joint smoothing distribution of  $\{\eta_t\}$  and  $\{\xi_t^P\}$  given  $\mathbf{Z}_1, \dots, \mathbf{Z}_T$ , the **posterior mean** of  $\mathbf{Y}_t^P$  at the set of locations  $P$  at time  $t$  is,

$$\begin{aligned}\mathbf{Y}_{t|T}^P &= \begin{pmatrix} \mathbf{Y}_{t|T}^{(1)P} \\ \mathbf{Y}_{t|T}^{(2)P} \end{pmatrix} \\ &= \mathbf{X}_t^P \alpha_t + \mathbf{S}_t^P \eta_{t|T} + \xi_{t|T}^P.\end{aligned}$$



## Prediction standard error matrix

The mean squared prediction error matrix (equivalently the posterior covariance matrix) can be calculated as:

$$\begin{aligned}\mathbf{M}_{t|T}^P &\equiv \mathbb{E} \left( \left[ \mathbf{Y}_t^P - \mathbf{Y}_{t|T}^P \right] \left[ \mathbf{Y}_t^P - \mathbf{Y}_{t|T}^P \right]' \right) \\ &= \begin{pmatrix} \mathbf{M}_{t|T}^{(1,1)P} & \mathbf{M}_{t|T}^{(1,2)P} \\ \mathbf{M}_{t|T}^{(2,1)P} & \mathbf{M}_{t|T}^{(2,2)P} \end{pmatrix} \\ &= \mathbf{S}_t^P \mathbf{P}_{t|T} \mathbf{S}_t^{P'} + \mathbf{R}_{t|T}^P + 2 \mathbf{S}_t^P \mathbf{W}_{t|T}^P,\end{aligned}$$

$$\text{where } \mathbf{M}_{t|T}^{(k,m)P} \equiv \mathbb{E} \left( \left[ \mathbf{Y}_t^{(k)P} - \mathbf{Y}_{t|T}^{(k)P} \right] \left[ \mathbf{Y}_t^{(m)P} - \mathbf{Y}_{t|T}^{(m)P} \right]' \right).$$



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Notes on STDF

STDF has attractive features that make it suitable for large remote sensing datasets,

- ▶ It is fast and scalable to large data inputs,
- ▶ It exploits the inter-process correlation for improved accuracy,
- ▶ It takes advantage of both *temporal* and *spatial* dependence in the data.



National Aeronautics and  
Space Administration

**Jet Propulsion Laboratory**  
California Institute of Technology  
Pasadena, California

## Outline

Introduction

Prelude: spatial (non-temporal) interpolation for massive datasets

Spatio-Temporal Data Fusion

Application to Lower Atmosphere CO<sub>2</sub>

Conclusion



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Lower-atmosphere CO<sub>2</sub>

- ▶ Deriving global distribution of lower-atmosphere CO<sub>2</sub> over time is important for studying 'sources' and 'sinks.'
- ▶ The Greenhouse gases Observing SATellite (GOSAT) provides total-column CO<sub>2</sub>, while the Atmospheric InfraRed Sounder (AIRS) provides mid-tropospheric CO<sub>2</sub>.
- ▶ We will derive joint predictions of total-column CO<sub>2</sub> and mid-tropospheric CO<sub>2</sub> and taking a (weighted) difference to obtain lower atmosphere CO<sub>2</sub>.



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Application outline

- ▶ We select GOSAT and AIRS data over the continental United States between June and August of 2009.
- ▶ We make joint-prediction of total-column CO<sub>2</sub> and mid-tropospheric CO<sub>2</sub>, and use weighted differencing to derive predictions of lower atmosphere CO<sub>2</sub>.
- ▶ Predictions of lower atmosphere CO<sub>2</sub> will be compared to coincident aircraft data from NOAA.
- ▶ We also compare the performance of STDF with an alternative interpolation methodology (locally weighted regression).



## Observing tracks and atmospheric sensitivity

### Example of satellite orbits and sensitivity




Figure: Example of GOSAT and AIRS sensitivity



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## ACOS and AIRS input data

- ▶ Within our domain, we have 3,869 ACOS data points and 40,564 AIRS data points.
- ▶ We group the data over these three months into 3-day blocks.
- ▶ For the elements of the vector of basis functions, we use local bisquare functions.
- ▶ The covariate function  $\mathbf{t}(\cdot)$  are defined using a constant 1, latitude, and longitude.
- ▶ Given the joint prediction,  $(\hat{Y}_{t|T,ACOS}(\mathbf{s}), \hat{Y}_{t|T,AIRS}(\mathbf{s}))'$ , we estimate lower atmosphere CO2 as a simple linear combination

$$\hat{Y}_{t|T,LA}(\mathbf{s}) = \frac{7}{5} \hat{Y}_{t|T,ACOS}(\mathbf{s}) + \frac{2}{5} \hat{Y}_{t|T,AIRS}(\mathbf{s}).$$






National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## ACOS and AIRS input example

### Example of input data





National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## STDF output example




Figure: STDF output

- Run time for entire 3-month period: 4 minutes on a 3.06 GHz machine with an Intel Duo Core processor.



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## NOAA lower-atmosphere CO<sub>2</sub> data

- ▶ The National Oceanic and Atmospheric Administration (NOAA) has been sampling lower-atmospheric CO<sub>2</sub> through a series of aircraft flights over Beaver Crossing, Nebraska and Lamont, Oklahoma.
- ▶ We can compare the NOAA aircraft data at these two locations against the corresponding 95% confidence intervals for STDF lower-atmospheric CO<sub>2</sub>.



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## NOAA lower-atmosphere CO<sub>2</sub> comparison




Figure: STDF outputs (red intervals) vs NOAA data (colored circles).




National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Comparison vs loess

- ▶ To compare against loess, we randomly 6 time blocks, and designate a small, fixed area as a reserve region.
- ▶ All data falling within the reserve region within those 6 time blocks are withheld as test data.
- ▶ We apply STDF and loess predict the value of ACOS and AIRS CO<sub>2</sub> at the test locations.






National Aeronautics and  
Space Administration


Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Comparison vs loess - results


ACOS residuals: Loess




AIRS residuals: Loess



ACOS residuals: STDF



AIRS residuals: STDF





National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Outline

### Introduction

Prelude: spatial (non-temporal) interpolation for massive datasets

Spatio-Temporal Data Fusion

Application to Lower Atmosphere CO<sub>2</sub>

### Conclusion



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Conclusions

- ▶ STDF is fast: feasible for large remote sensing datasets.
- ▶ Results look reasonable by comparison to aircraft data for this example.
- ▶ Applicable to other types of remote sensing data, e.g., aerosols, clouds, soil moisture...
- ▶ Extensions: Bayesian inference, application to remote sensing radiances, etc.
- ▶ Questions and/or comments: contact Hai Nguyen at [hai.nguyen@jpl.nasa.gov](mailto:hai.nguyen@jpl.nasa.gov).





National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Acknowledgments

The authors would like to thank Colm Sweeney of the NOAA Earth System Research Laboratory for providing the in-situ aircraft CO<sub>2</sub> measurements used in this study.

This research is supported by NASA's Earth Science Technology Office under its Advanced Information Technology Program.

This work was partially performed at the Jet Propulsion Laboratory, California Institution of Technology under contract with the National Aeronautics and Space Administration. Government sponsorship acknowledged. Copyright 2012, all rights reserved.



National Aeronautics and  
Space Administration

**Jet Propulsion Laboratory**  
California Institute of Technology  
Pasadena, California

# Appendix



Let  $\theta^{[b]}$  be the parameter vector at the  $b$ -th EM iteration. The conditional expectations and covariance matrices for the “missing data” are defined as:

$$\eta_{t|T}^{[b]} \equiv E_{\theta^{[b]}}(\eta_t | \mathbf{Z}_{1:T})$$

$$\xi_{t|T}^{[b]} \equiv E_{\xi^{[b]}}(\xi_t | \mathbf{Z}_{1:T})$$

$$\mathbf{P}_{t|T}^{[b]} \equiv \text{var}_{\theta^{[b]}}(\eta_t | \mathbf{Z}_{1:T})$$

$$\mathbf{R}_{t|T}^{[b]} \equiv \text{var}_{\theta^{[b]}}(\xi_t | \mathbf{Z}_{1:T})$$

$$\mathbf{W}_{t|T}^{[b]} \equiv \text{cov}_{\theta^{[b]}}(\eta_t, \xi_t | \mathbf{Z}_{1:T})$$

$$\mathbf{P}_{t,t-1|T}^{[b]} \equiv \text{cov}_{\theta^{[b]}}(\eta_t, \eta_{t-1} | \mathbf{Z}_{1:T}).$$



The cross-covariance term,  $\mathbf{P}_{t,t-1|T} \equiv \text{cov}(\boldsymbol{\eta}_t, \boldsymbol{\eta}_{t-1} | \mathbf{Z}_{1:T})$ , is given by,

$$\begin{aligned}\mathbf{P}_{T,T-1|T} &= (\mathbf{I}_r - \mathbf{P}_{T|T-1} \mathbf{S}'_T [\mathbf{S}_T \mathbf{P}_{T|T-1} \mathbf{S}'_T + \mathbf{D}_T]^{-1} \mathbf{S}_T) \\ &\quad \times \mathbf{H}_T \mathbf{P}_{T-1|T-1} \\ \mathbf{P}_{t,t-1|T} &= \mathbf{P}_{t|t} \mathbf{J}'_{t-1} + \mathbf{J}_t (\mathbf{P}_{t+1,t|T} - \mathbf{H}_{t+1} \mathbf{P}_{t|T}) \mathbf{J}'_{t-1},\end{aligned}$$

where  $\mathbf{I}_r$  is the  $r \times r$  identity matrix, and define,

$$\mathbf{L}_t^{[b+1]} \equiv \mathbf{P}_{t,t-1|T}^{[b]} + \boldsymbol{\eta}_t^{[b]} \boldsymbol{\eta}_{t-1|T}^{[b]'}.$$



The EM updates for  $\theta^{[b+1]}$  are:

$$\alpha_t^{[b+1]} = (\mathbf{X}_t' \mathbf{Q} \mathbf{V}_t^{-1} \mathbf{Q} \mathbf{X}_t)^{-1} \mathbf{X}_t' \mathbf{Q} \mathbf{V}_t^{-1} [\mathbf{Z}_t - \mathbf{S}_t \eta_{t|T}^{[b]} - \xi_{t|T}^{[b]}],$$

$$\mathbf{K}_0^{[b+1]} = \mathbf{P}_{0|T}^{[b]} + \eta_{0|T}^{[b]} \eta_{0|T}^{[b]'}.$$

$$(\sigma_{\xi,t}^{(1)})^2 [b+1] = \frac{1}{N_t^{(1)}} \text{trace} \left( \left( \mathbf{E}_t^{-1} [\mathbf{R}_{t|T}^{[b]} + \xi_{t|T}^{[b]} \xi_{t|T}^{[b]'}] \right)_{[1, N^{(1)}]} \right)$$

$$(\sigma_{\xi,t}^{(2)})^2 [b+1] = \frac{1}{N_t^{(2)}} \text{trace} \left( \left( \mathbf{E}_t^{-1} [\mathbf{R}_{t|T}^{[b]} + \xi_{t|T}^{[b]} \xi_{t|T}^{[b]'}] \right)_{[N^{(1)+1}, N]} \right)$$

$$\mathbf{H}^{[b+1]} = \left( \sum_{t=1}^T \mathbf{L}_t^{[b+1]} \right) \left( \sum_{t=0}^{T-1} \mathbf{K}_t^{[b+1]} \right)^{-1}$$

$$\mathbf{U}^{[b+1]} = \left( \sum_{t=1}^T \mathbf{K}_t^{[b+1]} - \mathbf{H}^{[b+1]} \sum_{t=1}^T \mathbf{L}_t^{[b+1]'} \right) / T.$$



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## EM estimation

Convergence of the parameter estimates may be monitored through the negative log-likelihood

$$-\log L(\theta) = \frac{1}{2} \sum_{t=1}^T \log |\Sigma_{\beta,t}(\theta)| + \frac{1}{2} \sum_{t=1}^T \beta_t' \Sigma_{\beta,t}(\theta)^{-1} \beta_t.$$



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Weighted differencing

We make the following simplifying assumptions,

- ▶ The air pressure at the surface of the Earth is 1000 hectopascals (hPa) and the air pressure at the satellite instrument is 0 hPa.
- ▶ The middle troposphere is the portion of the atmosphere between 500 hPa and 300 hPa.
- ▶ The CO<sub>2</sub> concentration above 300 hPa can be ignored.



National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Weighted differencing

Given total column CO<sub>2</sub>,  $Y_{\text{ACOS}}(\mathbf{s})$ , and mid-tropospheric CO<sub>2</sub>,  $Y_{\text{AIRS}}(\mathbf{s})$ , at a location  $\mathbf{s}$ , we approximated lower-atmospheric CO<sub>2</sub>,  $Y_{\text{LA}}(\mathbf{s})$ , as a simple linear combination,

$$\begin{aligned} Y_{\text{LA}}(\mathbf{s}) &= \frac{(1000 - 300) Y_{\text{ACOS}}(\mathbf{s}) - (500 - 300) Y_{\text{AIRS}}(\mathbf{s})}{1000 - 500} \\ &= \frac{7}{5} Y_{\text{ACOS}}(\mathbf{s}) - \frac{2}{5} Y_{\text{AIRS}}(\mathbf{s}). \end{aligned}$$





National Aeronautics and  
Space Administration

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

## Weighted differencing

From the weighted difference above, it is straightforward to obtain the prediction standard error at location  $\mathbf{s}$ ,

$$\sigma_{LA}^2(\mathbf{s}) \equiv \begin{pmatrix} 7/5, -2/5 \end{pmatrix} \mathbf{M}_{t|T}(\mathbf{s}) \begin{pmatrix} 7/5, -2/5 \end{pmatrix}',$$

where  $\mathbf{M}_{t|T}(\mathbf{s})$  is the prediction-error matrix for the CO<sub>2</sub> prediction vector  $\hat{\mathbf{Y}}_{t|T}(\mathbf{s}) \equiv (\hat{Y}_{t|T,ACOS}(\mathbf{s}), \hat{Y}_{t|T,AIRS}(\mathbf{s}))'$ .